

Effects of omega-3 fatty acid supplementation during pregnancy: A systematic review

Jaydeep Nag, A. Jothi Priya, R. Gayatri Devi

ABSTRACT

Omega-3 fatty acids are essential and can only be obtained from the diet. The requirements during pregnancy have not been established, but likely exceed that of a non-pregnant state. Omega-3 fatty acids are critical for fetal neurodevelopment and may be important for the timing of gestation and birth weight as well. Most pregnant women likely do not get enough omega-3 fatty acids because the major dietary source, seafood, is restricted to 2 servings a week. For pregnant women to obtain adequate omega-3 fatty acids, a variety of sources should be consumed: Vegetable oils, 2 low-mercury fish servings a week, and supplements (fish oil or algae-based docosahexaenoic acid).

KEY WORDS: Docosahexaenoic acid, Eicosapentaenoic acid, Fish oil, Omega-3 fatty acids, Pregnancy, Prenatal supplements

INTRODUCTION

In the field of perinatal nutrition, polyunsaturated fatty acids (PUFAs) of the omega-3 and omega-6 groups have gained recent attention because of their important functions in fetal and newborn neurodevelopment and because of their roles in inflammation.^[1-3]

Two PUFAs, arachidonic acid (AA) and docosahexaenoic acid (DHA), are critical to fetal and infant central nervous system (CNS) growth and development.^[1,4] Embedded in the cell membrane phospholipid, AA is involved in cell signaling pathways and cell division and serves as an inflammatory precursor for eicosanoids. The DHA concentration is high in retinal and brain membrane phospholipids, and it is involved in visual and neural function and neurotransmitter metabolism.^[2] During the last trimester, the fetus accrues about 50–70 mg a day of 1 omega-3 fatty acid, DHA.^[5,6] Both maternal DHA intake and circulating DHA concentrations are important determinants of fetal blood concentrations of DHA.^[3] Babies accrue DHA into the CNS up until about 18 months of age.^[7,8]

Although research into the specific pathways affected by these PUFAs is still in its infancy, there is enough

understanding to draw conclusions and make recommendations about their dietary intake during the perinatal period. For the purposes of this review, we focus primarily on the omega-3 fatty acids, DHA and eicosapentaenoic acid (EPA) and the omega-6 fatty acid, AA. We present information about how the absolute and relative concentrations of these fatty acids affect pregnancy outcome and fetal neurodevelopment.

Intake of omega-6-rich oils found in sunflower, corn, and cottonseed oils should be minimized because they are converted to substrates that compete with EPA. Pregnant women should reduce their intake of these oils and substitute others that are rich in omega-3 fatty acids such as flaxseed, canola, and soybean oil.

Of the 20 or so edible fatty acids, only omega-3 and omega-6 fatty acids cannot be synthesized by the body.^[9] All of the omega-3 and omega-6 fatty acids accumulated by the fetus must ultimately be derived from the mother by placental transfer. Omega refers to where the first double bond appears on the fatty acid chain from the carboxyl end. The typical American diet is replete with omega-6 fatty acids, particularly linoleic acid, which is readily converted to AA. Omega-6 fatty acids can be found in the vegetable oils used in processed foods, fried foods, and condiments such as salad dressings.^[10,11] One teaspoon of corn oil can satisfy the daily omega-6 requirement, but most individuals eat 10–20 times that amount.^[10,11]

Access this article online

Website: jrsolutions.info

ISSN: 0975-7619

Department of Physiology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India

*Corresponding author: Mrs. Jothi Priya, Department of Physiology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India, E-mail: a.jothipriya88@gmail.com

Received on: 13-02-2019; Revised on: 17-03-2019; Accepted on: 23-04-2019

In contrast, the intake of omega-3 fatty acids is suboptimal.^[9-11] The richest dietary sources of omega-3 fatty acids are from marine sources, fish oil supplements, and selected vegetable oils such as flaxseed (57% omega-3 fatty acids), canola (11% omega-3 fatty acids), and soybean (8% omega-3 fatty acids). Most individuals in the United States do not consume these omega-3-rich foods on a regular basis.^[12] The ratio of dietary omega-6/omega-3 fatty acids in the American diet approximates 10–25:1; the paleolithic diet was probably closer to 1:1.^[10]

The omega-3 fatty acid EPA and the omega-6 fatty acid AA are essential structural components of every cell in the body. Both EPA and AA serve as precursors for biologically active compounds called eicosanoids. These fatty acids compete for the enzyme systems cyclooxygenase, which makes prostaglandins (PGs) and thromboxanes (TXs) and lipoxygenase, which makes leukotrienes. Diets that are rich in omega-6 fatty acids produce potent eicosanoids, whereas a diet with a more balanced intake of omega-6 and omega-3 fatty acids makes less inflammatory and less immunosuppressive eicosanoids.^[13]

Specific to pregnancy, although both DHA and AA appear to be essential to fetal CNS development, the relatively poor intake of EPA coupled with the high intake of linoleic acid (which produces AA), may affect pregnancy outcome by altering the balance of the eicosanoids produced.^[2,5,6,14-16] A high ratio of AA to EPA may promote untoward effects such as pre-term labor and pre-eclampsia.^[2,17] A linoleic acid-rich diet produces an abundance of AA, which serves as a precursor of the potent 2-series PGs E_2 and $PGF_2\alpha$ and the vasoconstrictor TXA_2 . Both PGE_2 and $PGF_2\alpha$ are closely associated with the initiation of labor and preterm labor, whereas TXA_2 has been associated with pre-eclampsia.^[18,19] Whereas the omega-3 fatty acid, DHA, is not usually considered to be involved with eicosanoid formation, EPA is a precursor for the 3-series of PGs and produces PGE_3 and PGI_3 , which promote relaxation of myometrium.^[20,21] Furthermore, EPA and DHA competitively displace AA in the membrane phospholipids and thereby reduce the production of 2-series eicosanoids.^[22,23] Thus, a diet that provides a closer balance of omega-3 to omega-6 fatty acids may be as important to pregnant women as the absolute individual plasma levels of these fatty acids.

Nonetheless, in pregnancy, the real significance of EPA may be related to its role in mediating DHA and AA concentrations across the placenta rather than its production of the relatively less potent PGI_3 . Free fatty acids need to be bound for transfer to the fetal circulation. Selective transport across the human placenta for individual fatty acids has been suggested

as 1 mechanism to explain greater concentrations of some PUFAs such as DHA and AA in the fetal, rather than maternal, circulation. In this regard, the effect of EPA on mRNA expression of fatty acid transport proteins (FATPs) becomes important.^[24] EPA, but not DHA, has been positively correlated with mRNA expression of all membrane proteins. Thus, higher maternal EPA concentrations may increase FATP expression (FATP-4 in particular) that, in turn, has been shown to increase cord blood DHA levels.^[24] In addition, the free fatty acids need to be bound to fatty acid binding proteins (FABPs) to gain entry into placental and fetal cells. Because, as noted above, EPA has been positively correlated with mRNA expression of all membrane proteins, higher EPA concentrations also lead to increase expression of FABPs including B-FABP, which is strongly expressed in developing brain cells and has a strong affinity for DHA.^[25]

Because only about 4%–11% of DHA is retroconverted to EPA,^[10,11,26] pregnant women who just take DHA supplements, without any dietary EPA, may be unable to produce the right balance of eicosanoids and may limit the transport and uptake of DHA into fetal cells.^[10,11,24-26]

CLINICAL TRIALS

Benefits of Omega-3 Fatty Acids

Much of the interest in omega-3 fatty acid intake and pregnancy began in the early 1980s when Danish investigators determined that women living on the Faroe Islands delivered babies that were 194 g heavier and had gestation lengths 4 days longer than babies born in Denmark.^[21] The Faroese diet had substantially more omega-3 fatty acids and less omega-6 fatty acids than a Danish diet. Red blood cell (RBC) fatty acid content (expressed as the ratio of omega-3 to omega-6) was significantly higher in the Faroese pregnant women than in Danish pregnant women.^[27] From this study, the authors theorized that a Danish woman who adjusted her diet to increase her blood ratio of these fatty acids by 20% would expect to increase gestation 5.7 days.^[27]

Following up on their earlier epidemiologic work, the same Danish group then randomized pregnant women at week 30 of gestation to either a fish oil supplement (2.7 g omega-3 fatty acids, of which 920 mg were DHA), olive oil, or no supplement.^[20] Women in the fish oil group had a gestational period 4 days longer than women taking either olive oil or no supplement. The babies born to the women in the fish oil group also weighed 107 g more than babies born to the women in the olive oil group and 43 g more than those born to those mothers without supplementation. The supplemental amount of dietary omega-3 fatty acids used in this study was larger than a typical Danish

diet, which only provided about 10% of this amount (i.e., 270 mg of omega-3 fatty acids). Thus, it appeared that rather high amounts of omega-3 fatty acids need to be consumed to affect gestation and fetal weight.

In a different Scandinavian trial, Norwegian pregnant women ($n = 341$) between gestational weeks 17 and 19 were randomized to consume omega-3 fatty acids from cod liver oil (i.e., 2632 mg omega-3 fatty acids with 1183 mg as DHA) or corn oil daily, until 3 months after delivery.^[28] Unlike the previous study, the primary outcomes of gestational length and birth weight did not differ between the two groups.^[20] Interestingly, however, they did note that infants with the highest quartile concentration of DHA in the cord blood had a gestational period 9.3 days longer than those in the lowest quartile. Thus, it appeared that although increased delivery of DHA to the fetus may be of benefit, how much DHA gets to the fetus may be entirely related to maternal intake.

Later, these same investigators evaluated the fatty acid content of the blood of the mothers and babies from the previous study.^[28,29] No maternal blood fatty acid differences occurred at baseline. However, at the end of pregnancy, the maternal plasma omega-3 fatty acids (e.g., EPA and DHA) and the ratio of omega-3 to omega-6 fatty acids were higher in the mothers who were assigned to the cod liver oil supplement compared with those in the corn oil group. In addition, fetal DHA umbilical cord plasma phospholipids were 23% higher in babies born to mothers in the cod liver oil group compared with those born to mothers in the corn oil group. Thus, healthy pregnant women taking about 1 g of DHA a day from a marine oil supplement were able to deliver substantially more DHA to the fetus than those who were not given marine oils. Finally, and most importantly, 4 years later, when these authors evaluated this same Norwegian cohort of infants, they concluded that the children had higher mental processing scores when born to mothers supplemented with cod liver oil (rich in EPA and DHA, omega-3 PUFAs) during pregnancy and lactation, compared with children of mothers who were supplemented with corn oil (rich in linoleic acid, omega-6 PUFAs).^[30]

In the same vein, an Australian group also looked at infant cognition in a similar trial and found similar results.^[30,31] At 20 weeks of gestation until delivery, 98 women were randomized to receive either fish oil (2.2 g DHA and 1.1 g EPA/day) or olive oil. At birth, infants from mothers in the fish oil-supplemented group had higher DHA and EPA levels and lower AA levels in the cord blood compared with children born to mothers in the olive oil-supplemented group. When evaluated at 2.5 years of age, children in the fish oil-supplemented group ($n = 33$) had higher scores for

eye and hand coordination compared with those in the olive oil group. In this study, as in others, eye and hand coordination scores and other tests of mental acuity correlated with omega-3 levels (EPA and DHA) in cord blood erythrocytes and inversely correlated with omega-6 levels (AA).^[23,31]

BENEFITS OF DHA-ONLY SUPPLEMENTATION

The time of the most rapid neural and retinal development occurs in the second half of pregnancy, mainly during the third trimester. On this basis, supplementation of the maternal diet later in pregnancy with omega-3 fatty acids, especially DHA, was thought to be especially important.^[3,7,11,14,15] In a trial focused on the benefits of naturally occurring DHA from food, a sample of pregnant Inuit women living in Arctic Quebec was assessed.^[23] Maternal DHA blood concentrations were shown to be directly related to cord plasma phospholipid levels, and the correlation of the DHA/AA ratio in maternal and cord plasma was even stronger. In this study, higher DHA cord blood concentration was related to longer gestation, better visual acuity, and mental and psychomotor skills at 6 months and 11 months, suggesting that raising DHA concentrations alone may independently yield some benefits. (It must be noted that the Inuit population has a dietary intake of EPA and DHA that is higher than most other nations because of their regular intake of fish and marine animals. Such foods are also rich in EPA, but it was not measured).

A second study provides large amounts of DHA as a supplement explored its effects during pregnancy and, later, on fetal cognition.^[3] From week 16 of gestation until delivery, healthy Canadian women (average age, 33 years) who were not taking supplemental fish oil capsules were randomized to 400 mg of algae-derived DHA or a blend of corn and soybean oil. The purpose of the study was to compare RBC DHA levels, dietary intakes of DHA and other omega-3 fatty acids, and infant visual acuity at 60 days of age. At week 36, DHA in the supplemented women was 32% higher than the control group, and no differences were found between the groups for EPA and AA, both of which decreased. Infants were 3 times more likely to have low visual acuity scores in the placebo group than the DHA group, indicating that the usual diet may have had insufficient DHA to maximize visual acuity in infants. In fact, omega-3 fatty acid deficiency, as measured by RBC DHA, was more prevalent in the control group than in the DHA-supplemented group.

LACK OF OMEGA-3 FATTY ACIDS IN THE WESTERN DIET

The trials outlined above (and numerous others not mentioned) provide a compelling argument for the

essential nature of omega-3 PUFAs during pregnancy. Yet, the typical Western diet is notably deficient with respect to them. Three studies demonstrated that pregnant women likely consume inadequate omega-3 fatty acids.^[7,32,33] In a Canadian study, pregnant women were found to consume only 1.5 g of omega-3 fatty acids per day, of which only 117 mg were from EPA and DHA.^[7] This is below what is considered adequate for a non-pregnant woman and quite inadequate in DHA.

A Danish survey study conducted in high-risk pregnant women showed that dietary omega-3 fatty acids were inversely related to the risk of preterm delivery.^[33] Women consuming <150 mg of omega-3 fatty acids a day (i.e., <0.5 oz of fish per day) were at the highest risk of delivering prematurely. Of those who ate no seafood and did not use fish oil supplements, the rate of preterm delivery was 7.1%, compared with 1.9% in those who regularly ate fish. Thus, for high-risk pregnant women, omega-3 fatty acid intake seemed to have an important effect on reducing spontaneous premature births. Unlike the Canadian study, the Danish women who regularly ate fish appeared to obtain adequate omega-3 fatty acids from the diet alone and did not require supplementation.

Finally, in the United States, there is considerable evidence of inadequate seafood consumption. In a review of the 1999–2000 National Health and Nutrition Examination Survey, Smith and Sahyoun concluded that only 19% of American adults consume the recommended 2 servings of fish each week.^[34,35,36,37] Furthermore, Oken *et al.* reported that after a well-publicized January 2001 federal advisory regarding methylmercury contamination in certain fish species, seafood consumption of all species decreased in pregnant American women, even as recommendations were publicized by nutritional groups as to the benefits of seafood for pregnant women.^[38,39]

The mercury found in 2 servings of the approved seafood poses little risk to the fetus.^[40,41] The only evidence of fetal neurodevelopmental abnormalities was seen in infants whose mothers consumed known, high amounts of mercury-rich seafood. The mercury intake of these women was much higher than what an American woman would get from eating 2 servings of low-mercury fish per week (10–30 ppm vs. <0.05 ppb, respectively). These conclusions are further corroborated by the project viva data where children born to mothers eating more than 2 weekly servings of fish performed better on language and visual and motor tests at 3 years of age compared with children born to mothers who ate less than this amount.^[40] Only 12% of the 341 pregnant women consumed more than 2 servings of fish per week. No adverse effects of maternal mercury

concentrations were observed. Maternal consumption of fish lower in mercury resulted in stronger benefits to the offspring and thus underscores the importance of consuming low-mercury seafood during pregnancy.

For those seeking to avoid seafood, there are few non-supplement options. Plant-based omega-3 fatty acids (i.e., α -linolenic acid [ALA]), such as flaxseed oil, are poorly converted to the biologically active omega-3 fatty acid EPA and convert even less to DHA.^[3,10,11,26,43] The range of conversion of ALA to EPA is generally between 0.2% and 9% although some authors have suggested that women of childbearing age may be able to convert up to 21% of their dietary ALA to EPA. Regardless of which conversion factor is correct, trying to obtain all omega-3 fatty acids from plant-based oils requires ingestion of too many fat calories.

Unfortunately, it is impossible for pregnant women to meet their omega-3 fatty acid requirements from omega-3-rich vegetable oils and 2 servings of seafood a week. Two servings of fish per week only provide about 100–250 mg/day of omega-3 fatty acids, of which 50–100 mg is from DHA; plant-based oils supply insignificant amounts of EPA and no DHA. During pregnancy, the dietary goal for omega-3 fatty acids is 650 mg, of which 300 is DHA.^[44] Thus, to make up the omega-3 fatty acid deficit in the diet, pregnant women are left with essentially 2 choices: Fish oil supplements supplying EPA and DHA or algae-derived DHA. Specifically, depending on the omega-3 content of the seafood consumed during the week, each day pregnant women would need an additional 400–550 mg of omega-3 PUFAs (EPA and DHA), of which about 225 mg should be DHA.

Fish oil supplements are commercially available from multiple companies. The amount of EPA and DHA per capsule varies but most contain one-third to one-half of these omega-3 fatty acids (e.g., in a 1000-mg capsule, 300 or 500 mg would come from EPA and DHA). Most commercially available fish oil supplements contain < 1–2 ppb of mercury compared with <0.05 ppb in seafood, but because the amount of fish oil used (1–2 g a day) is much smaller than what is consumed when eating seafood (200 g twice a week), the risk of mercury toxicity is inconsequential. Cod liver oil is considered a less desirable source of EPA and DHA because it is also rich in Vitamin A, unlike other fish oils. Long-term use of Vitamin A has been associated with an increased risk of osteoporosis.^[45]

Meaningful vegetarian sources of DHA are essentially limited to algae-derived DHA from Martek biosciences (Columbia, MD). Using a strain of algae, *Cryptocodinium cohnii*, which is a naturally high producer of DHA, DHA oil is produced in the US food and drug administration-inspected, environmentally

controlled manufacturing facilities. Although free of any environmental contaminants such as mercury, the oils do not contain any EPA, and data demonstrating the benefits in pregnancy of DHA alone are lacking.

Recommendations

There is little doubt that pregnant women need at least as many omega-3 fatty acids as non-pregnant women and likely need more DHA.^[10,14,44] The recommendations for dietary omega-3 fatty acids should be adopted at the onset of pregnancy, but there may be benefits for all women who are considering becoming pregnant. Given concerns for mercury toxicity with overconsumption of certain fish, to meet these recommendations, pregnant women will need to consume omega-3 fatty acids from 3 sources: Vegetable oils, 2 servings of seafood per week, and omega-3 fatty acid supplements containing EPA and DHA or DHA alone. Intake of omega-6-rich oils found in sunflower, corn, and cottonseed oils should be minimized because they are converted to substrates that compete with EPA. Pregnant women should reduce their intake of these oils and substitute others that are rich in omega-3 fatty acids such as flaxseed, canola, and soybean oil.^[44,46]

CONCLUSION

Hence, this article is done to create an awareness about the omega-3 supplementation during pregnancy.

REFERENCES

- Birch EE, Garfield S, Castañeda Y, Hughbanks-Wheaton D, Uauy R, Hoffman D, *et al.* Visual acuity and cognitive outcomes at 4 years of age in a double-blind, randomized trial of long-chain polyunsaturated fatty acid-supplemented infant formula. *Early Hum Dev* 2007;83:279-84.
- Elias SL, Innis SM. Infant plasma trans, n-6, and n-3 fatty acids and conjugated linoleic acids are related to maternal plasma fatty acids, length of gestation, and birth weight and length. *Am J Clin Nutr* 2001;73:807-14.
- Innis SM, Friesen RW. Essential n-3 fatty acids in pregnant women and early visual acuity maturation in term infants. *Am J Clin Nutr* 2008;87:548-57.
- Birch EE, Castañeda YS, Wheaton DH, Birch DG, Uauy RD, Hoffman DR, *et al.* Visual maturation of term infants fed long-chain polyunsaturated fatty acid-supplemented or control formula for 12 mo. *Am J Clin Nutr* 2005;81:871-9.
- Clandinin MT, Chappell JE, Heim T, Swyer PR, Chance GW. Fatty acid utilization in perinatal de novo synthesis of tissues. *Early Hum Dev* 1981;5:355-66.
- Innis SM. Essential fatty acid transfer and fetal development. *Placenta* 2005;26:S70-5.
- Denomme J, Stark KD, Holub BJ. Directly quantitated dietary (n-3) fatty acid intakes of pregnant Canadian women are lower than current dietary recommendations. *J Nutr* 2005;135:206-11.
- Szajewska H, Horvath A, Koletzko B. Effect of n-3 long-chain polyunsaturated fatty acid supplementation of women with low-risk pregnancies on pregnancy outcomes and growth measures at birth: A meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2006;83:1337-44.
- Bell SJ, Bradley D, Forse RA, Bistrrian BR. The new dietary fats in health and disease. *J Am Diet Assoc* 1997;97:280-6.
- Kris-Etherton PM, Taylor DS, Yu-Poth S, Huth P, Moriarty K, Fishell V, *et al.* Polyunsaturated fatty acids in the food chain in the United States. *Am J Clin Nutr* 2000;71:179S-88.
- Burdge G. Alpha-linolenic acid metabolism in men and women: Nutritional and biological implications. *Curr Opin Clin Nutr Metab Care* 2004;7:137-44.
- CanolaInfo. Canola Oil Information Web Site. Available from: <http://www.canolainfo.org/health/index.php?page=12>. [Last accessed on 2008 Mar 20].
- Endres S, Ghorbani R, Kelley VE, Georgilis K, Lonnemann G, van der Meer JW, *et al.* The effect of dietary supplementation with n-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med* 1989;320:265-71.
- Jensen CL. Effects of n-3 fatty acids during pregnancy and lactation. *Am J Clin Nutr* 2006;83:1452S-7.
- Koletzko B, Larqué E, Demmelmair H. Placental transfer of long-chain polyunsaturated fatty acids (LC-PUFA). *J Perinat Med* 2007;35 Suppl 1:S5-11.
- Allen KG, Harris MA. The role of n-3 fatty acids in gestation and parturition. *Exp Biol Med (Maywood)* 2001;226:498-506.
- Wang Y, Walsh SW, Kay HH. Placental tissue levels of nonesterified polyunsaturated fatty acids in normal and preeclamptic pregnancies. *Hypertens Pregnancy* 2005;24:235-45.
- Olson DM. The role of prostaglandins in the initiation of parturition. *Best Pract Res Clin Obstet Gynaecol* 2003;17:717-30.
- Malatyalioglu E, Adam B, Yanik FF, Kökçü A, Alvrur M. Levels of stable metabolites of prostacyclin and thromboxane A2 and their ratio in normotensive and preeclamptic pregnant women during the antepartum and postpartum periods. *J Matern Fetal Med* 2000;9:173-7.
- Olsen SF, Sørensen JD, Secher NJ, Hedegaard M, Henriksen TB, Hansen HS, *et al.* Randomised controlled trial of effect of fish-oil supplementation on pregnancy duration. *Lancet* 1992;339:1003-7.
- Olsen SF, Hansen HS, Sørensen TI, Jensen B, Secher NJ, Sommer S, *et al.* Intake of marine fat, rich in (n-3)-polyunsaturated fatty acids, may increase birthweight by prolonging gestation. *Lancet* 1986;2:367-9.
- Smuts CM, Huang M, Mundy D, Plasse T, Major S, Carlson SE, *et al.* A randomized trial of docosahexaenoic acid supplementation during the third trimester of pregnancy. *Obstet Gynecol* 2003;101:469-79.
- Jacobson JL, Jacobson SW, Muckle G, Kaplan-Estrin M, Ayotte P, Dewailly E, *et al.* Beneficial effects of a polyunsaturated fatty acid on infant development: Evidence from the inuit of arctic Quebec. *J Pediatr* 2008;152:356-64.
- Larqué E, Krauss-Etschmann S, Campoy C, Hartl D, Linde J, Klingler M, *et al.* Docosahexaenoic acid supply in pregnancy affects placental expression of fatty acid transport proteins. *Am J Clin Nutr* 2006;84:853-61.
- Xu LZ, Sánchez R, Sali A, Heintz N. Ligand specificity of brain lipid-binding protein. *J Biol Chem* 1996;271:24711-9.
- Conquer JA, Holub BJ. Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects. *J Nutr* 1996;126:3032-9.
- Olsen SF, Hansen HS, Sommer S, Jensen B, Sørensen TI, Secher NJ, *et al.* Gestational age in relation to marine n-3 fatty acids in maternal erythrocytes: A study of women in the Faroe Islands and Denmark. *Am J Obstet Gynecol* 1991;164:1203-9.
- Helland IB, Saugstad OD, Smith L, Saarem K, Solvoll K, Ganes T, *et al.* Similar effects on infants of n-3 and n-6 fatty acids supplementation to pregnant and lactating women. *Pediatrics* 2001;108:E82.
- Helland IB, Saugstad OD, Saarem K, Van Houwelingen AC, Nylander G, Drevon CA, *et al.* Supplementation of n-3 fatty acids during pregnancy and lactation reduces maternal plasma lipid levels and provides DHA to the infants. *J Matern Fetal Neonatal Med* 2006;19:397-406.
- Helland IB, Smith L, Saarem K, Saugstad OD, Drevon CA.

- Maternal supplementation with very-long-chain n-3 fatty acids during pregnancy and lactation augments children's IQ at 4 years of age. *Pediatrics* 2003;111:e39-44.
31. Dunstan JA, Simmer K, Dixon G, Prescott SL. Cognitive assessment of children at age 2(1/2) years after maternal fish oil supplementation in pregnancy: A randomised controlled trial. *Arch Dis Child Fetal Neonatal Ed* 2008;93:F45-50.
 32. Oken E, Kleinman KP, Olsen SF, Rich-Edwards JW, Gillman MW. Associations of seafood and elongated n-3 fatty acid intake with fetal growth and length of gestation: Results from a US pregnancy cohort. *Am J Epidemiol* 2004;160:774-83.
 33. Olsen SF, Secher NJ. Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: Prospective cohort study. *BMJ* 2002;324:447-50.
 34. Smith KM, Sahyoun NR. Fish consumption: Recommendations versus advisories, can they be reconciled? *Nutr Rev* 2005;63:39-46.
 35. US Environmental Protection Agency Office of Water, Authors. Proceedings of the 2007 National Forum on Contaminants in Fish. Washington, DC: United States Environmental Protection Agency Office of Water; 2007. Available from: <http://www.epa.gov/waterscience/fish/forum/2007/pdf/section2e.pdf>. [Last accessed on 2008 Feb 28].
 36. US Department of Health and Human Services and US Environmental Protection Agency, Authors. Backgrounder for the 2004 FDA/EPA Consumer Advisory: What You Need to Know About Mercury in Fish and Shellfish. Washington, DC: US Environmental Protection Agency; 2004. Available from: <http://www.epa.gov/waterscience/fishadvice/factsheet.html>. [Last accessed on 2008 Feb 28].
 37. Nettleton JA. Omega-3 fatty acids: Comparison of plant and seafood sources in human nutrition. *J Am Diet Assoc* 1991;91:331-7.
 38. Oken E, Kleinman KP, Berland WE, Simon SR, Rich-Edwards JW, Gillman MW, *et al.* Decline in fish consumption among pregnant women after a national mercury advisory. *Obstet Gynecol* 2003;102:346-51.
 39. ACOG Updates Definitive Guide to Pregnancy, Authors. Your Pregnancy and Birth, Fourth Edition Gives Women the Latest News on Genetic Disorders, Prenatal Tests, DOs and DON'Ts, and Exercise. The American College of Obstetricians and Gynecologists Office of Communications; May 10; 2005. Available from: http://www.acog.org/from_home/publications/press_releases/nr05-10-05-2.cfm. [Last accessed on 2008 Feb 28].
 40. Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: Evaluating the risks and the benefits. *JAMA* 2006;296:1885-99.
 41. Mahaffey KR. Fish and shellfish as dietary sources of methylmercury and the omega-3 fatty acids, eicosahexaenoic acid and docosahexaenoic acid: Risks and benefits. *Environ Res* 2004;95:414-28.
 42. Oken E, Radesky JS, Wright RO, Bellinger DC, Amarasiwardena CJ, Kleinman KP, *et al.* Maternal fish intake during pregnancy, blood mercury levels, and child cognition at age 3 years in a US cohort. *Am J Epidemiol* 2008;167:1171-81.
 43. Pawlosky RJ, Hibbeln JR, Novotny JA, Salem N Jr. Physiological compartmental analysis of alpha-linolenic acid metabolism in adult humans. *J Lipid Res* 2001;42:1257-65.
 44. Simopoulos AP, Leaf A, Salem N Jr. Essentiality of and recommended dietary intakes for omega-6 and omega-3 fatty acids. *Ann Nutr Metab* 1999;43:127-30.
 45. Feskanich D, Singh V, Willett WC, Colditz GA. Vitamin A intake and hip fractures among postmenopausal women. *JAMA* 2002;287:47-54.
 46. Koletzko B, Cetin I, Brenna JT, Perinatal Lipid Intake Working Group, Child Health Foundation, Diabetic Pregnancy Study Group, *et al.* Dietary fat intakes for pregnant and lactating women. *Br J Nutr* 2007;98:873-7.

Source of support: Nil; Conflict of interest: None Declared